

# Matched Asymptotic Expansions of the Eigenvalues of a 3-D boundary-value problem relative to two cavities linked by a hole of small size

M'Barek Fares<sup>1</sup> Abdelkader Tizaoui<sup>2</sup> and Sébastien Tordeux<sup>2,\*</sup>

<sup>1</sup> *Electromagnetism and Acoustics, CERFACS, 42 Avenue Gaspard Coriolis, F-31100 Toulouse.*

<sup>2</sup> *Toulouse University, INSA-Toulouse, Mathematical Institute of Toulouse (UMR-CNRS 5219), 135 avenue de Rangueil, F-31077 Toulouse, France.*

---

**Abstract.** In this article, we consider a domain consisting of two cavities linked by a hole of small size. We derive a numerical method to compute an approximation of the eigenvalues of an elliptic operator without refining in the neighborhood of the hole. Several convergence rates are obtained and illustrated by numerical simulations.

**AMS subject classifications:** 34E05, 35J05, 65M60, 78M30, 78M35

**Key words:** Elliptic Operator, Matched Asymptotic Expansions, Eigenvalue Problem, Finite Elements.

---

## 1 Introduction

### 1.1 Motivation

In a lot of physical problems, the boundary of the computational domain is perforated. This configuration can lead to numerical difficulties when the diameter of the holes are really smaller than the other characteristic lengths. Indeed, it can be very costly to compute a sharp numerical approximation of the solution of such problems for two main reasons: With a standard method like finite elements or finite differences, a refined mesh cannot be avoided in the neighborhood of the hole; the mesh generation of a perforated structure can be a hard task.

Many authors have studied the effect of the perforation of the boundaries both from the theoretical and the numerical point of views, see for example [13, 16–19]. However, fewer results have been obtained for the eigenvalue problem in the case of a three dimensional domain.

---

\*Corresponding author. *Email address:* stordeux@insa-toulouse.fr

In [10], Gadyl'shin considered a two dimensional domain consisting of two domains linked by a small hole. He derived a complete asymptotic expansion of the scattering frequencies of the Laplacian operator equipped with Dirichlet boundary condition. In [2], these results were extended to the eigenvalues and eigenvectors of an elliptic operator with varying coefficients. In this paper, we are interested in a three dimensional configuration with varying coefficients and Neumann boundary condition.

## 1.2 A Neumann eigenvalue problem

**The geometry.** Let  $\Omega_{\text{int}}$  and  $\Omega_{\text{ext}}$  be two open subsets of  $\mathbb{R}^3$  with

$$\Omega_{\text{int}} \cap \Omega_{\text{ext}} = \emptyset \quad \text{and} \quad \exists \delta_0 > 0: ]-2\delta_0, 2\delta_0[^3 \cap \partial\Omega_{\text{int}} \cap \partial\Omega_{\text{ext}} = (]-2\delta_0, 2\delta_0[^2 \times \{0\}). \quad (1.1)$$

Let  $\Sigma \subset ]-1, 1[^2$  be an open subset of  $\mathbb{R}^2$ . For  $\delta < \delta_0$ , we consider the domain  $\Omega^\delta$ , see Figure 1, consisting of  $\Omega_{\text{ext}}$  and  $\Omega_{\text{int}}$  linked by an iris  $\Sigma_\delta = \delta\Sigma = \{(x, y) \in \mathbb{R}^2 : (\frac{x}{\delta}, \frac{y}{\delta}) \in \Sigma\}$

$$\Omega^\delta := \Omega_{\text{int}} \cup \Omega_{\text{ext}} \cup (\Sigma_\delta \times \{0\}) \subset \mathbb{R}^3. \quad (1.2)$$

This domain tends to  $\Omega := \Omega_{\text{int}} \cup \Omega_{\text{ext}} \subset \mathbb{R}^2$  when  $\delta \rightarrow 0$ .

**The original problem.** In these domains we consider the eigenvalue problems




Figure 1: The computational domain  $\Omega^\delta$ .

$$\left\{ \begin{array}{l} \text{Find } (\lambda_n^\delta, u_n^\delta) \in \mathbb{R} \times H^1(\Omega^\delta): \\ -\nabla \cdot (a \nabla u_n^\delta) = \lambda_n^\delta b u_n^\delta \text{ in } \Omega^\delta, \\ \partial_{\mathbf{n}} u_n^\delta = 0 \text{ on } \partial\Omega^\delta, \end{array} \right. \quad (1.3)$$

$$\left\{ \begin{array}{l} \text{Find } (\lambda_n, u_n) \in \mathbb{R} \times H^1(\Omega): \\ -\nabla \cdot (a \nabla u_n) = \lambda_n b u_n \text{ in } \Omega, \\ \partial_{\mathbf{n}} u_n = 0 \text{ on } \partial\Omega, \end{array} \right. \quad (1.4)$$

with  $a \in L^\infty(\Omega)$  and  $b \in L^\infty(\Omega)$  two functions of  $\Omega$  with  $\inf_{\mathbf{x} \in \Omega} a(\mathbf{x}) > 0$ ,  $\inf_{\mathbf{x} \in \Omega} b(\mathbf{x}) > 0$  and whose restrictions to  $\Omega_{\text{int}}$  and  $\Omega_{\text{ext}}$  are regular and can be expanded in the neighborhood of  $\mathbf{0}$  with the form

$$a|_{\Omega_{\text{ext}}}(\mathbf{x}) = \sum_{i,j,k \geq 0} a_{i,j,k}^{\text{ext}} x^i y^j z^k, \quad b|_{\Omega_{\text{ext}}}(\mathbf{x}) = \sum_{i,j,k \geq 0} b_{i,j,k}^{\text{ext}} x^i y^j z^k \quad \text{with } a_{i,j,k}^{\text{ext}}, b_{i,j,k}^{\text{ext}} \in \mathbb{R}, \quad (1.5)$$

$$a|_{\Omega_{\text{int}}}(\mathbf{x}) = \sum_{i,j,k \geq 0} a_{i,j,k}^{\text{int}} x^i y^j z^k, \quad b|_{\Omega_{\text{int}}}(\mathbf{x}) = \sum_{i,j,k \geq 0} b_{i,j,k}^{\text{int}} x^i y^j z^k \quad \text{with } a_{i,j,k}^{\text{int}}, b_{i,j,k}^{\text{int}} \in \mathbb{R}. \quad (1.6)$$

In order to shorten the expressions, we adopt the notations  $a_0 = a_{0,0,0}$  and  $b_0 = b_{0,0,0}$ . The discrete sets of eigenmodes  $(u_n^\delta, \lambda_n^\delta)_{n \geq 0}$  (resp.  $(u_n, \lambda_n)_{n \geq 0}$ ) can be chosen to be a bi-orthogonal basis of  $L^2(\Omega^\delta)$  and  $H^1(\Omega^\delta)$  (resp.  $L^2(\Omega)$  and  $H^1(\Omega)$ ) and to satisfy

$$0 = \lambda_0^\delta \leq \lambda_1^\delta \leq \lambda_2^\delta \leq \dots \quad \text{and} \quad \lim_{n \rightarrow +\infty} \lambda_n^\delta = +\infty, \quad (1.7)$$

respectively

$$0 = \lambda_0 = \lambda_1 \leq \lambda_2 \leq \dots \quad \text{and} \quad \lim_{n \rightarrow +\infty} \lambda_n = +\infty \quad \text{and} \quad \forall n \in \mathbb{N} \quad u_n|_{\Omega_{\text{int}}} = 0 \text{ or } u_n|_{\Omega_{\text{ext}}} = 0. \quad (1.8)$$

Some natural questions arise: Does the eigenvalue  $\lambda_n^\delta$  converge to  $\lambda_n$ ? Is it possible to obtain an asymptotic expansion of  $\lambda_n^\delta$ ? With this asymptotic expansion, is it possible to compute a numerical approximation of  $\lambda_n^\delta$  with a small computation cost?

### 1.3 Main theorem

The next Theorem gives positive answers to these three questions.

**Theorem 1.1.** *Let  $n \in \mathbb{N}$  and let  $\alpha$  be the positive real defined in Section 3.1.*

(i) *If  $\lambda_n$  is a simple eigenvalue of the limit problem, then  $\lambda_n^\delta$  can be expanded as follows*

$$\lambda_n^\delta = \lambda_n + 2\pi\alpha \frac{a_0^{\text{int}} a_0^{\text{ext}}}{a_0^{\text{int}} + a_0^{\text{ext}}} \frac{(\bar{u}_n(\mathbf{0}))^2}{\int_{\Omega} b(u_n)^2} \delta + O_{\delta \rightarrow 0}(\delta^2 \ln \delta), \quad (1.9)$$

with the notation

$$\bar{u}(\mathbf{0}) = u|_{\Omega_{\text{ext}}}(\mathbf{0}) \quad \text{if } u|_{\Omega_{\text{int}}} = 0 \quad \text{and} \quad \bar{u}(\mathbf{0}) = u|_{\Omega_{\text{int}}}(\mathbf{0}) \quad \text{if } u|_{\Omega_{\text{ext}}} = 0. \quad (1.10)$$

(ii) If  $\lambda_n$  is a double eigenvalue of the limit problem ( $\lambda_n = \lambda_{n+1}$ ), then  $\lambda_n^\delta = \lambda_n + O_{\delta \rightarrow 0}(\delta^2)$  and  $\lambda_{n+1}^\delta$  can be expanded as follows

$$\lambda_{n+1}^\delta = \lambda_n + 2\pi\alpha \frac{a_0^{int} a_0^{ext}}{a_0^{int} + a_0^{ext}} \left( \frac{(\bar{u}_n(\mathbf{0}))^2}{\int_{\Omega} (bu_n)^2} + \frac{(\bar{u}_{n+1}(\mathbf{0}))^2}{\int_{\Omega} (bu_{n+1})^2} \right) \delta + O_{\delta \rightarrow 0}(\delta^2 \ln \delta). \quad (1.11)$$

**Remark 1.1.** Formulas (1.9) and (1.11) involve only quantities which are independent of  $\delta$ . Consequently truncating these expressions by eliminating the remainders furnish an approximation of  $\lambda_n^\delta$  which does not require any mesh refinement.

**Remark 1.2.** Due to (1.8), one has either  $u_n|_{\Omega_{int}} = 0$  or  $u_n|_{\Omega_{ext}} = 0$ . Consequently, (1.10) always defines  $\bar{u}(\mathbf{0})$ .

#### 1.4 Matched asymptotic expansions

The first order asymptotic expansions of an eigenvalue  $\lambda_n^\delta$  reads

$$\lambda_n^\delta = \lambda_n^0 + \delta \lambda_n^1 + o_{\delta \rightarrow 0}(\delta) \quad (1.12)$$

and has been derived in parallel to the derivation of the first order asymptotic expansion of the eigenfunction  $u_n^\delta$ . The model (1.3) involves two characteristic lengths of different magnitude: the size of the hole  $\delta$  which is much smaller than the diameter of the cavity. Multiple scalings should be used to obtain an approximation of the eigenvector  $u_n^\delta$  uniformly valid. The first scaling corresponds to the  $\mathbf{x}$ -variable and takes care of the cavity phenomena. The second scaling  $\mathbf{X} = \mathbf{x}/\delta$  permits to describe the boundary layer phenomena located in the neighborhood of the hole. Guided by the well known method of Matching of Asymptotic Expansions, see [11] and [20], we look for the asymptotic expansions of the two functions  $\delta \mapsto u_n^\delta(\mathbf{x})$  and  $\delta \mapsto \Pi_n^\delta(\mathbf{X}) := u_n^\delta(\delta \mathbf{X})$ . At first order, they take the form

$$u_n^\delta(\mathbf{x}) = u_n^0(\mathbf{x}) + \delta u_n^1(\mathbf{x}) + o_{\delta \rightarrow 0}(\delta), \quad (1.13)$$

$$\Pi_n^\delta(\mathbf{X}) := u_n^\delta(\delta \mathbf{X}) = \Pi_n^0(\mathbf{X}) + \delta \Pi_n^{1,0}(\mathbf{X}) + \delta \ln \delta \Pi_n^{1,1}(\mathbf{X}) + o_{\delta \rightarrow 0}(\delta). \quad (1.14)$$

The functions  $u_n^i$  are defined on the domain  $\Omega$  and are possibly singular in the neighborhood of the origin. The functions  $\Pi_n^i$  are defined on a normalized version of the neighborhood of the hole

$$\widehat{\Omega} := \mathbb{R}^3 \setminus \{(x, y, 0) : (x, y) \in \mathbb{R}^2 \setminus \Sigma\}. \quad (1.15)$$

These two asymptotic expansions match asymptotically in an intermediate region.

**Remark 1.3.** The presence of poly-logarithmic gauge functions  $\delta^u \ln^p \delta$  is rather not classical for three dimensional problems. They are due to the non constancy of the coefficient

*a.* Indeed in the case of a constant  $a$  we have  $\Pi^{1,1} = 0$ . Moreover in the case of non constant coefficient, it is possible to show that Logarithms appear also in the eigenvalue and far-field expansions of second order which take the form

$$\lambda_n^\delta = \lambda_n^0 + \delta \lambda_n^1 + \delta^2 \lambda_n^{2,0} + \delta^2 \ln \delta \lambda_n^{2,1} + \underset{\delta \rightarrow 0}{o}(\delta^2), \quad (1.16)$$

$$u_n^\delta(\mathbf{x}) = u_n^0(\mathbf{x}) + \delta u_n^1(\mathbf{x}) + \delta^2 u_n^{2,0}(\mathbf{x}) + \delta^2 \ln \delta u_n^{2,1}(\mathbf{x}) + \underset{\delta \rightarrow 0}{o}(\delta^2). \quad (1.17)$$

## 1.5 Content

In this paper we will not give the complete proof of Theorem 1.1 which is mainly based on the third order matched asymptotic expansions, on the min-max principle [14] and on a quasi-mode approach [8]. A complete proof for a two dimensional Dirichlet-Laplacian can be found in [3].

This paper will be focused on the description of the coefficients of the asymptotic expansions (1.12), (1.13) and (1.14) and on the question of their existence and uniqueness in the case of a simple eigenvalue. Moreover, the formulas (1.9) and (1.11) will be shown to be in good agreement with some direct numerical simulations.

## 2 Matched asymptotic expansions of simple eigenvalues

In this section, we suppose that  $\lambda_n$  is a simple eigenvalue. We will only deal with the case  $u_n|_{\Omega_{\text{ext}}} = 0$ . The case  $u_n|_{\Omega_{\text{int}}} = 0$  can be deduced by symmetry.

The coefficients of the three asymptotic expansions (1.12), (1.13) and (1.14) have been formally derived using the Van Dyke matching principle [20]. The problems defining these coefficients will be proved to be well-posed in Section 3.

### 2.1 The limit coefficients

The far-field and eigenvalue limit coefficients are rather naturally defined by

$$u_n^0 = u_n \text{ and } \lambda_n^0 = \lambda_n. \quad (2.1)$$

Moreover, the near-field limit coefficient  $\Pi_n^0$  is defined on  $\widehat{\Omega}$  by

$$\begin{cases} -\nabla \cdot (a_0 \nabla \Pi_n^0) = 0 \text{ in } \widehat{\Omega}, & \partial_{\mathbf{n}} \Pi_n^0 = 0 \text{ on } \partial \widehat{\Omega}, \\ \Pi_n^0|_{\widehat{\Omega}_{\text{int}}}(\mathbf{X}) = u_n^0|_{\Omega_{\text{int}}}(\mathbf{0}) + \underset{R \rightarrow +\infty}{o}(1) \text{ and } \Pi_n^0|_{\widehat{\Omega}_{\text{ext}}}(\mathbf{X}) = \underset{R \rightarrow +\infty}{o}(1), \end{cases} \quad (2.2)$$

with  $R = \sqrt{X^2 + Y^2 + Z^2}$ ,  $\widehat{\Omega}_{\text{int}}$  the lower half space and  $\widehat{\Omega}_{\text{ext}}$  the upper half space. As it will be proved in Section 3.1, see Remark 3.1 and Lemma 3.1, this coefficient can be expanded

in the neighborhood of  $+\infty$  with the form

$$\begin{cases} \Pi_n^0|_{\widehat{\Omega}_{\text{int}}}(\mathbf{X}) = u_n^0|_{\Omega_{\text{int}}}(\mathbf{0}) - \frac{a_0^{\text{ext}} \alpha}{a_0^{\text{int}} + a_0^{\text{ext}}} u_n|_{\Omega_{\text{int}}}(\mathbf{0}) \frac{1}{R} + \underset{R \rightarrow +\infty}{O}\left(\frac{1}{R^2}\right), \\ \Pi_n^0|_{\widehat{\Omega}_{\text{ext}}}(\mathbf{X}) = \frac{a_0^{\text{int}} \alpha}{a_0^{\text{int}} + a_0^{\text{ext}}} u_n|_{\Omega_{\text{int}}}(\mathbf{0}) \frac{1}{R} + \underset{R \rightarrow +\infty}{O}\left(\frac{1}{R^2}\right). \end{cases} \quad (2.3)$$

## 2.2 The first order coefficients

The first order asymptotic expansions are given by (1.12), (1.13) and (1.14) where the first order coefficients  $\lambda_n^1 \in \mathbb{R}$ ,  $u_n^1 : \Omega \rightarrow \mathbb{R}$ ,  $\Pi_n^{1,0} : \widehat{\Omega} \rightarrow \mathbb{R}$  and  $\Pi_n^{1,1} : \widehat{\Omega} \rightarrow \mathbb{R}$  remain to be defined. The terms  $u_n^1$  and  $\lambda_n^1$  are solutions of the well-posed coupled problem

$$\begin{cases} \nabla \cdot (a \nabla u_n^1) + \lambda_n b u_n^1 = -\lambda_n^1 b u_n \text{ in } \Omega, \quad \partial_{\mathbf{n}} u_n^1 = 0 \text{ on } \partial\Omega \setminus \{\mathbf{0}\}, \\ u_n^1 + \frac{a_0^{\text{ext}} \alpha}{a_0^{\text{int}} + a_0^{\text{ext}}} u_n|_{\Omega_{\text{int}}}(\mathbf{0}) \frac{1}{r} \in H^1(\Omega_{\text{int}}), \\ u_n^1 - \frac{a_0^{\text{int}} \alpha}{a_0^{\text{int}} + a_0^{\text{ext}}} u_n|_{\Omega_{\text{int}}}(\mathbf{0}) \frac{1}{r} \in H^1(\Omega_{\text{ext}}), \end{cases} \quad (2.4)$$

with  $r = \sqrt{x^2 + y^2 + z^2}$ . The coefficient  $u_n^1$  can then be expanded in the neighborhood of  $\mathbf{0}$  with the help of the Kondratiev's theory, see [7, 12],

$$\begin{cases} u_n^1|_{\Omega_{\text{int}}}(\mathbf{x}) = -\frac{a_0^{\text{ext}} \alpha}{a_0^{\text{int}} + a_0^{\text{ext}}} u_n|_{\Omega_{\text{int}}}(\mathbf{0}) \frac{1}{r} + s_n^1(\mathbf{x}) + r_n^1(\mathbf{x}), \\ u_n^1|_{\Omega_{\text{ext}}}(\mathbf{x}) = \frac{a_0^{\text{int}} \alpha}{a_0^{\text{int}} + a_0^{\text{ext}}} u_n|_{\Omega_{\text{int}}}(\mathbf{0}) \frac{1}{r} + s_n^1(\mathbf{x}) + r_n^1(\mathbf{x}), \end{cases} \quad (2.5)$$

with

$$\begin{aligned} s_n^1|_{\Omega_{\text{int}}}(\mathbf{x}) &= \frac{a_0^{\text{ext}} \alpha}{a_0^{\text{int}} + a_0^{\text{ext}}} \frac{u_n|_{\Omega_{\text{int}}}(\mathbf{0})}{2} \left( \frac{a_{1,0,0}^{\text{int}}}{a_0^{\text{int}}} \frac{x}{r} + \frac{a_{0,1,0}^{\text{int}}}{a_0^{\text{int}}} \frac{y}{r} + \frac{a_{0,0,1}^{\text{int}}}{a_0^{\text{int}}} \left( \frac{z}{r} + \ln \frac{r-z}{2} \right) \right), \\ s_n^1|_{\Omega_{\text{ext}}}(\mathbf{x}) &= -\frac{a_0^{\text{int}} \alpha}{a_0^{\text{int}} + a_0^{\text{ext}}} \frac{u_n|_{\Omega_{\text{int}}}(\mathbf{0})}{2} \left( \frac{a_{1,0,0}^{\text{ext}}}{a_0^{\text{ext}}} \frac{x}{r} + \frac{a_{0,1,0}^{\text{ext}}}{a_0^{\text{ext}}} \frac{y}{r} + \frac{a_{0,0,1}^{\text{ext}}}{a_0^{\text{ext}}} \left( \frac{z}{r} - \ln \frac{r+z}{2} \right) \right) \end{aligned}$$

and  $r_n^1 : \Omega \rightarrow \mathbb{R}$  such that the two restrictions  $r_n^1|_{\Omega_{\text{int}}}$  and  $r_n^1|_{\Omega_{\text{ext}}}$  are continuous in the neighborhood of  $\mathbf{0}$ .

The coefficients  $\Pi_n^{1,0}$  and  $\Pi_n^{1,1}$  are defined in the infinite domain  $\widehat{\Omega}$  by

$$\begin{cases} -\nabla \cdot (a_0 \nabla \Pi_n^{1,0}) = \nabla \cdot \left( (a_{1,0,0} X + a_{0,1,0} Y + a_{0,0,1} Z) \nabla \Pi_n^0 \right) \text{ in } \widehat{\Omega}, \quad \partial_{\mathbf{n}} \Pi_n^{1,0} = 0 \text{ on } \partial\widehat{\Omega}, \\ \Pi_n^{1,0}|_{\widehat{\Omega}_{\text{int}}}(\mathbf{X}) = \partial_x u_n^0|_{\Omega_{\text{int}}}(\mathbf{0}) X + \partial_y u_n^0|_{\Omega_{\text{int}}}(\mathbf{0}) Y + s_n^1|_{\Omega_{\text{int}}}(\mathbf{X}) + r_n^1|_{\Omega_{\text{int}}}(\mathbf{0}) + \underset{r \rightarrow 0}{o}(1), \\ \Pi_n^{1,0}|_{\widehat{\Omega}_{\text{ext}}}(\mathbf{X}) = s_n^1|_{\Omega_{\text{ext}}}(\mathbf{X}) + r_n^1|_{\Omega_{\text{ext}}}(\mathbf{0}) + \underset{r \rightarrow 0}{o}(1), \end{cases} \quad (2.6)$$

$$\begin{cases} -\nabla \cdot (a_0 \nabla \Pi_n^{1,1}) = 0 \text{ in } \widehat{\Omega}, & \partial_{\mathbf{n}} \Pi_n^{1,1} = 0 \text{ on } \partial \widehat{\Omega}, \\ \Pi_n^{1,1}|_{\widehat{\Omega}_{\text{int}}}(\mathbf{X}) = \frac{u_n|_{\Omega_{\text{int}}}(\mathbf{0})}{2} \frac{a_0^{\text{ext}} \alpha}{a_0^{\text{int}} + a_0^{\text{ext}}} \frac{a_{0,0,1}^{\text{int}}}{a_0^{\text{int}}} + o_{r \rightarrow 0}(1), \\ \Pi_n^{1,1}|_{\widehat{\Omega}_{\text{ext}}}(\mathbf{X}) = \frac{u_n|_{\Omega_{\text{int}}}(\mathbf{0})}{2} \frac{a_0^{\text{int}} \alpha}{a_0^{\text{int}} + a_0^{\text{ext}}} \frac{a_{0,0,1}^{\text{ext}}}{a_0^{\text{ext}}} + o_{r \rightarrow 0}(1). \end{cases} \quad (2.7)$$

### 3 Existence of the coefficients of the asymptotic expansions

#### 3.1 Existence and uniqueness of the $\Pi_n^i$

The question of existence and uniqueness of the Laplacian problems equipped with Neumann boundary condition is a rather well understood topic, see for example [1]. For  $F \in L^2(\widehat{\Omega})$  compactly supported, let us recall that there exists a unique  $\Pi \in H_{\text{loc}}^1(\widehat{\Omega})$  satisfying

$$\nabla \cdot (a_0 \nabla \Pi) = F \text{ in } \widehat{\Omega}, \quad \partial_{\mathbf{n}} \Pi = 0 \text{ on } \partial \widehat{\Omega}, \quad \Pi = o_{R \rightarrow +\infty}(1). \quad (3.1)$$

This is mainly due to the Hardy inequality

$$\exists \gamma > 0: \quad \gamma \left( \|\nabla \Pi\|_{L^2(\widehat{\Omega})} + \left\| \frac{\Pi}{1+R} \right\|_{L^2(\widehat{\Omega})} \right) \leq \|\nabla \Pi\|_{L^2(\widehat{\Omega})}, \quad \forall \Pi \in K_0^1 \quad (3.2)$$

and to the equivalence of the last problem with the variational formulation

$$\text{Find } \Pi \in K_0^1: \int_{\widehat{\Omega}} a_0 \nabla \Pi \cdot \nabla \Pi' = \int_{\widehat{\Omega}} F \Pi' \text{ for all } \Pi' \in K_0^1 \quad (3.3)$$

with the Kondratiev's space

$$K_0^1 := \left\{ \Pi: \widehat{\Omega} \rightarrow \mathbb{R}: \nabla \Pi \in L^2(\widehat{\Omega}) \text{ and } \frac{\Pi}{1+R} \in L^2(\widehat{\Omega}) \right\}. \quad (3.4)$$

We will now prove the existence and uniqueness of  $\Pi^0$  and  $\Pi^{1,1}$ . Let  $\Psi_{\text{int}}$  and  $\Psi_{\text{ext}}$  be two regular cut-off functions satisfying

$$\begin{cases} \Psi_{\text{int}}(\mathbf{X}) = 0 \text{ in } \widehat{\Omega}_{\text{ext}}, \\ \Psi_{\text{int}}(\mathbf{X}) = \varphi(R) \text{ in } \widehat{\Omega}_{\text{int}}, \end{cases} \quad (3.5) \quad \begin{cases} \Psi_{\text{ext}}(\mathbf{X}) = 0 \text{ in } \widehat{\Omega}_{\text{int}}, \\ \Psi_{\text{ext}}(\mathbf{X}) = \varphi(R) \text{ in } \widehat{\Omega}_{\text{ext}}, \end{cases} \quad (3.6)$$

with  $\varphi(R) = 0$  for  $R < 1$  and  $\varphi(R) = 1$  for  $R > 2$ .

**Theorem 3.1.** *For all reals  $A$  and  $B$ , there exists a unique  $\Pi_{A,B} \in H_{\text{loc}}^1(\widehat{\Omega})$  satisfying*

$$\begin{cases} -\nabla \cdot (a_0 \nabla \Pi_{A,B}) = 0 \text{ in } \widehat{\Omega}, & \partial_{\mathbf{n}} \Pi_{A,B} = 0 \text{ on } \partial \widehat{\Omega}, \\ \Pi_{A,B}|_{\widehat{\Omega}_{\text{int}}}(\mathbf{X}) = A + o_{R \rightarrow +\infty}(1) \text{ and } \Pi_{A,B}|_{\widehat{\Omega}_{\text{ext}}}(\mathbf{X}) = B + o_{R \rightarrow +\infty}(1). \end{cases} \quad (3.7)$$

*Proof.* We consider the function  $\Pi(\mathbf{X}) = \Pi_{A,B}(\mathbf{X}) - \Psi_{\text{int}}(\mathbf{X}) A - \Psi_{\text{ext}}(\mathbf{X}) B$  which satisfies (3.1) with

$$F(\mathbf{X}) = -\nabla \cdot \left( a_0 \left( A \nabla \Psi_{\text{int}}(\mathbf{X}) + B \nabla \Psi_{\text{ext}}(\mathbf{X}) \right) \right). \quad (3.8)$$

Since the function  $F$  is compactly supported, the problem (3.7) is well posed.  $\square$

**Remark 3.1.** The coefficients  $\Pi_n^0$  and  $\Pi_n^{1,1}$ , defined by (2.2) and (2.6), can be expressed as

$$\begin{aligned} \Pi_n^0 &= \Pi_{A,B} \text{ with } A = u_n|_{\Omega_{\text{int}}}(\mathbf{0}) \text{ and } B = 0, \\ \Pi_n^{1,1} &= \Pi_{A,B} \text{ with } A = \frac{1}{2} \frac{a_0^{\text{ext}} \alpha}{a_0^{\text{int}} + a_0^{\text{ext}}} \frac{a_{0,0,1}^{\text{int}}}{a_0^{\text{int}}} u_n|_{\Omega_{\text{int}}}(\mathbf{0}) \text{ and } B = \frac{1}{2} \frac{a_0^{\text{int}} \alpha}{a_0^{\text{int}} + a_0^{\text{ext}}} \frac{a_{0,0,1}^{\text{ext}}}{a_0^{\text{ext}}} u_n|_{\Omega_{\text{int}}}(\mathbf{0}). \end{aligned}$$

**Remark 3.2.** The existence and uniqueness of  $\Pi^{1,0}$  can be as well demonstrated. This requires extra arguments that are not central in our study. We have chosen not to give the details in this article.

Now we will be interested in the obtention of the asymptotic expansion of  $\Pi_{A,B}$  in the neighborhood of  $R = +\infty$ . Let us introduce the function  $\Pi_\star \in H_{\text{loc}}^1(\widehat{\Omega})$  satisfying

$$\begin{cases} \nabla \cdot (a_0 \nabla \Pi_\star) = 0 \text{ in } \widehat{\Omega}, & \partial_{\mathbf{n}} \Pi_\star = 0 \text{ on } \partial \widehat{\Omega}, \\ \Pi_\star|_{\widehat{\Omega}_{\text{int}}} = 1 + \underset{R \rightarrow +\infty}{o}(1) \text{ and } \Pi_\star|_{\widehat{\Omega}_{\text{ext}}} = \underset{R \rightarrow +\infty}{o}(1). \end{cases} \quad (3.9)$$

The function  $\Pi_\star$  is related to  $\Pi_{A,B}$  by  $\Pi_{A,B} = B + (A - B)\Pi_\star$ . In order to solve this problem, we will use a simple layer formulation based on the operator  $\mathbf{S}$

$$\mathbf{S}: (H^{\frac{1}{2}}(\Sigma))^* \mapsto H^{\frac{1}{2}}(\Sigma), \quad \lambda \mapsto \mathbf{S}\lambda(\mathbf{X}) = \frac{1}{4\pi} \int_{\Sigma} \frac{\lambda(\mathbf{X}')}{\|\mathbf{X} - \mathbf{X}'\|} dX'. \quad (3.10)$$

Taking into account the two transmission conditions  $\Pi_\star|_{\Sigma_{\text{int}}} = \Pi_\star|_{\Sigma_{\text{ext}}}$  and  $a_0^{\text{int}} \partial_z \Pi_\star|_{\Sigma_{\text{int}}} = a_0^{\text{ext}} \partial_z \Pi_\star|_{\Sigma_{\text{ext}}}$  and the representation formulas, see [9],

$$\Pi_\star|_{\Sigma_{\text{int}}} = 1 + 2\mathbf{S}\partial_z \Pi_\star|_{\Sigma_{\text{int}}} \quad \text{and} \quad \Pi_\star|_{\Sigma_{\text{ext}}} = -2\mathbf{S}\partial_z \Pi_\star|_{\Sigma_{\text{ext}}}, \quad (3.11)$$

we get the formulation

$$\text{Find } \lambda_\star \in (H^{\frac{1}{2}}(\Sigma))^* \text{ such that } \mathbf{S}\lambda_\star = 1 \text{ on } \Sigma, \quad (3.12)$$

with  $\lambda_\star$  related to the two normal derivatives by

$$\partial_z \Pi_\star|_{\Sigma_{\text{int}}} = -\frac{a_0^{\text{ext}}}{a_0^{\text{ext}} + a_0^{\text{int}}} \frac{\lambda_\star}{2} \quad \text{and} \quad \partial_z \Pi_\star|_{\Sigma_{\text{ext}}} = -\frac{a_0^{\text{int}}}{a_0^{\text{int}} + a_0^{\text{ext}}} \frac{\lambda_\star}{2}. \quad (3.13)$$



Then, the restrictions of  $\Pi_*$  to  $\widehat{\Omega}_{\text{int}}$  and  $\widehat{\Omega}_{\text{ext}}$  are given by the representation formula

$$\Pi_* = \begin{cases} 1 - \frac{a_0^{\text{ext}}}{a_0^{\text{ext}} + a_0^{\text{int}}} \left( \frac{1}{4\pi} \int_{\Sigma} \frac{\lambda_*(\mathbf{X}')}{\|\mathbf{X} - \mathbf{X}'\|} d\mathbf{X}' \right) & \text{in } \widehat{\Omega}_{\text{int}}, \\ + \frac{a_0^{\text{int}}}{a_0^{\text{ext}} + a_0^{\text{int}}} \left( \frac{1}{4\pi} \int_{\Sigma} \frac{\lambda_*(\mathbf{X}')}{\|\mathbf{X} - \mathbf{X}'\|} d\mathbf{X}' \right) & \text{in } \widehat{\Omega}_{\text{ext}}. \end{cases} \quad (3.14)$$

Expanding (3.14), the first order asymptotic expansion of  $\Pi_*$  for  $R = \|\mathbf{X}\| \rightarrow +\infty$  reads

$$\Pi_* = \begin{cases} 1 - \frac{a_0^{\text{ext}}}{a_0^{\text{ext}} + a_0^{\text{int}}} \frac{\alpha}{R} + O_{R \rightarrow +\infty} \left( \frac{1}{R^2} \right) & \text{in } \widehat{\Omega}_{\text{int}}, \\ + \frac{a_0^{\text{int}}}{a_0^{\text{ext}} + a_0^{\text{int}}} \frac{\alpha}{R} + O_{R \rightarrow +\infty} \left( \frac{1}{R^2} \right) & \text{in } \widehat{\Omega}_{\text{ext}}, \end{cases} \quad (3.15)$$

where  $\alpha = \frac{1}{4\pi} \int_{\Sigma} \lambda_*(\mathbf{X}') d\mathbf{X}'$  with  $\lambda_*$  defined by (3.12). The next Lemma follows.

**Lemma 3.1.**  $\Pi_{A,B}$  can be expanded with the form

$$\begin{cases} \Pi_{A,B}|_{\widehat{\Omega}_{\text{int}}}(\mathbf{X}) = A + (B - A) \frac{a_0^{\text{ext}}}{a_0^{\text{int}} + a_0^{\text{ext}}} \frac{\alpha}{R} + O_{R \rightarrow +\infty} \left( \frac{1}{R^2} \right), \\ \Pi_{A,B}|_{\widehat{\Omega}_{\text{ext}}}(\mathbf{X}) = B + (A - B) \frac{a_0^{\text{int}}}{a_0^{\text{int}} + a_0^{\text{ext}}} \frac{\alpha}{R} + O_{R \rightarrow +\infty} \left( \frac{1}{R^2} \right). \end{cases} \quad (3.16)$$

**Remark 3.3.** The coefficient  $\alpha$  depends on the shape of  $\Sigma$  but is independent of the value of  $a_0^{\text{int}}$  and  $a_0^{\text{ext}}$ . This quantity is related to the so called acoustic conductivity of the hole  $c$  and to the effective size of the hole  $s$  by  $s = \frac{c}{\pi} = \alpha\delta$ , see [19]. It can be numerically computed for every  $\Sigma$ , see Section 4. Moreover it has been analytically computed for some simple  $\Sigma$  with area  $\mathcal{A}$ , see [15]

- $\alpha = \frac{2\rho}{\pi} = \frac{2}{\pi} \sqrt{\frac{\mathcal{A}}{\pi}}$  for  $\Sigma$  a circle with radius  $\rho$ ,
- $\alpha = \frac{2}{\pi} \sqrt{\frac{\mathcal{A}}{\pi} \frac{(\pi/2)\sqrt{b/a}}{K(1-b^2/a^2)}}$  for  $\Sigma$  an ellipse with minor axes  $a$  and  $b$ . The function  $K$  denotes the complete elliptic integral of first kind.

The coefficient  $\alpha$  can be rather easily approximated by the coefficient  $\alpha$  of the circle with same area

$$\alpha_{\text{app}} = \frac{2}{\pi} \sqrt{\frac{\mathcal{A}}{\pi}} \quad (3.17)$$

This approximation does not require any numerical computation and is rather accurate for not too elongated holes. In figure 2 we illustrate the accuracy of this approximation in the case of an ellipse of minor axes  $a$  and  $b$  and a rectangle  $[0, a] \times [0, b]$ . The relative error is less than 5% for  $0.4 < b/a < 1$ .




Figure 2: Efficiency of the approximation of  $\alpha$  by  $\alpha_{app} = \frac{\pi}{2} \sqrt{\frac{A}{\pi}}$  for ellipses (left) and for rectangles (right).

### 3.2 Existence and uniqueness of $u_n^1$ and $\lambda_n^1$

The two coefficients  $u_n^1$  and  $\lambda_n^1$  do have to solve the problem (2.4). The following Lemma ensures the existence and uniqueness of  $u_n^1$  and  $\lambda_n^1$  up to the knowledge of the  $u_n$ -component of  $u_n^1$ . This component can be chosen arbitrarily.

**Lemma 3.2.** *Problem (2.4) has solutions. Moreover if  $(u_n^1, \lambda_n^1)$  and  $(u_{n,*}^1, \lambda_{n,*}^1)$  are solutions, one has  $\lambda_n^1 = \lambda_{n,*}^1$  and*

$$\lambda_n^1 = 2\pi\alpha \frac{a_0^{\text{int}} a_0^{\text{ext}}}{a_0^{\text{int}} + a_0^{\text{ext}}} \frac{(u_n|_{\Omega_{\text{int}}}(\mathbf{0}))^2}{\int_{\Omega_{\text{int}}} b(u_n)^2} \quad \text{and} \quad \exists \gamma \in \mathbb{R} : u_{n,*}^1 - u_n^1 = \gamma u_n. \quad (3.18)$$

*Proof.* • The function  $u_n^1$  does not belong to  $H^1(\Omega)$ . Consequently, the Fredholm alternative cannot be directly applied. For this reason, we introduce the auxiliary function  $\omega_n^1 \in H^1(\Omega)$

$$\begin{cases} \omega_n^1|_{\Omega_{\text{int}}}(\mathbf{x}) = u_n^1(\mathbf{x}) + \chi(r) u_n|_{\Omega_{\text{int}}}(\mathbf{0}) \frac{a_0^{\text{ext}}}{a_0^{\text{int}} + a_0^{\text{ext}}} \frac{\alpha}{r}, \\ \omega_n^1|_{\Omega_{\text{ext}}}(\mathbf{x}) = u_n^1(\mathbf{x}) - \chi(r) u_n|_{\Omega_{\text{int}}}(\mathbf{0}) \frac{a_0^{\text{int}}}{a_0^{\text{int}} + a_0^{\text{ext}}} \frac{\alpha}{r}, \end{cases} \quad (3.19)$$

with  $\chi$  a regular cut-off function satisfying (see (1.1))

$$\chi(z) = 1 \quad \text{if } z \leq \delta_0, \quad \text{and} \quad \chi(z) = 0 \quad \text{if } z \geq 2\delta_0. \quad (3.20)$$

Using (2.4),  $\omega_n^1$  belongs to  $H^1(\Omega)$  and satisfies

$$\nabla \cdot (a \nabla \omega_n^1) + \lambda_n b \omega_n^1 = F_n^1 \quad \text{in } \Omega \quad \text{and} \quad \partial_n \omega_n^1 = 0 \quad \text{on } \partial\Omega, \quad (3.21)$$

with  $F_n^1 \in (H^1(\Omega))^*$  defined by

$$\begin{cases} F_n^1|_{\Omega_{\text{int}}}(\mathbf{x}) = -\lambda_n^1 b u_n(\mathbf{x}) + \left( \nabla \cdot (a \nabla) + \lambda_n b \right) \left( \chi(r) u_n|_{\Omega_{\text{int}}}(\mathbf{0}) \frac{a_0^{\text{ext}}}{a_0^{\text{int}} + a_0^{\text{ext}}} \frac{\alpha}{r} \right), \\ F_n^1|_{\Omega_{\text{ext}}}(\mathbf{x}) = -\left( \nabla \cdot (a \nabla) + \lambda_n b \right) \left( \chi(r) u_n|_{\Omega_{\text{int}}}(\mathbf{0}) \frac{a_0^{\text{ext}}}{a_0^{\text{int}} + a_0^{\text{ext}}} \frac{\alpha}{r} \right). \end{cases}$$

Since  $\lambda_n^0$  is a simple eigenvalue of the elliptic operator on  $\Omega_{\text{int}}$ , the problem (3.21) defining  $\omega_n^1$  has solutions if and only if

$$\int_{\Omega_{\text{int}}} F_n^1 u_n = 0. \quad (3.22)$$

Moreover this solution is determined up to its  $u_n$ -component, ie. if  $\omega_n^1$  and  $\omega_n^{1,*}$  are two solutions of (3.21) then

$$\exists \gamma \in \mathbb{R}: \quad \omega_n^1 = \omega_n^{1,*} + \gamma u_n. \quad (3.23)$$

- The necessary and sufficient condition (3.22) for existence of  $u_n^1$  will now be made explicit. Since  $u_n = 0$  in  $\Omega_{\text{ext}}$  and  $\nabla \cdot (a \nabla u_n) + \lambda_n b u_n = 0$  in  $\Omega_{\text{int}}$ , (3.22) takes the form

$$\begin{aligned} \lambda_n^1 \int_{\Omega_{\text{int}}} b (u_n)^2 &= \int_{\Omega_{\text{int}}} \nabla \cdot \left( a \nabla \left( \chi(r) u_n|_{\Omega_{\text{int}}}(\mathbf{0}) \frac{a_0^{\text{ext}}}{a_0^{\text{int}} + a_0^{\text{ext}}} \frac{\alpha}{r} \right) \right) u_n \\ &\quad - \int_{\Omega_{\text{int}}} \chi(r) u_n|_{\Omega_{\text{int}}}(\mathbf{0}) \frac{a_0^{\text{ext}}}{a_0^{\text{int}} + a_0^{\text{ext}}} \frac{\alpha}{r} (\nabla \cdot (a \nabla u_n)). \end{aligned} \quad (3.24)$$

Let us denote by  $B_\eta$  the ball of center  $\mathbf{0}$  and of radius  $\eta$ . Since the domain  $\Omega_{\text{int}} \setminus B_\eta$  tends to  $\Omega_{\text{int}}$  when  $\eta \rightarrow 0$ , we have due to Lebesgues Theorem

$$\begin{aligned} \lambda_n^1 \int_{\Omega_{\text{int}}} b (u_n)^2 &= \lim_{\eta \rightarrow 0^+} \left[ \int_{\Omega_{\text{int}} \setminus B_\eta} \nabla \cdot \left( a \nabla \left( \chi(r) u_n|_{\Omega_{\text{int}}}(\mathbf{0}) \frac{a_0^{\text{ext}}}{a_0^{\text{int}} + a_0^{\text{ext}}} \frac{\alpha}{r} \right) \right) u_n \right. \\ &\quad \left. - \int_{\Omega_{\text{int}} \setminus B_\eta} \chi(r) u_n|_{\Omega_{\text{int}}}(\mathbf{0}) \frac{a_0^{\text{ext}}}{a_0^{\text{int}} + a_0^{\text{ext}}} \frac{\alpha}{r} (\nabla \cdot (a \nabla u_n)) \right]. \end{aligned} \quad (3.25)$$

Two Green formulas lead to

$$\begin{aligned} &\int_{\Omega_{\text{int}} \setminus B_\eta} \nabla \cdot \left( a \nabla \left( \chi(r) u_n|_{\Omega_{\text{int}}}(\mathbf{0}) \frac{a_0^{\text{ext}}}{a_0^{\text{int}} + a_0^{\text{ext}}} \frac{\alpha}{r} \right) \right) u_n \\ &= - \left[ \int_{\frac{\pi}{2}}^{\pi} \int_0^{2\pi} a \partial_r \left( \chi(r) u_n|_{\Omega_{\text{int}}}(\mathbf{0}) \frac{a_0^{\text{ext}}}{a_0^{\text{int}} + a_0^{\text{ext}}} \frac{\alpha}{r} \right) u_n r^2 \sin(\theta) d\theta d\varphi \right] (r = \eta) \\ &\quad - \int_{\Omega_{\text{int}} \setminus B_\eta} a \nabla \left( \chi(r) u_n|_{\Omega_{\text{int}}}(\mathbf{0}) \frac{a_0^{\text{ext}}}{a_0^{\text{int}} + a_0^{\text{ext}}} \frac{\alpha}{r} \right) \nabla u_n r^2 \sin(\theta) dr d\theta d\varphi, \end{aligned} \quad (3.26)$$

$$\begin{aligned}
& \int_{\Omega_{\text{int}} \setminus B_\eta} \chi(r) u_n |_{\Omega_{\text{int}}}(\mathbf{0}) \frac{a_0^{\text{ext}}}{a_0^{\text{int}} + a_0^{\text{ext}}} \frac{\alpha}{r} (\nabla \cdot (a \nabla u_n)) \\
&= - \left[ \int_{\frac{\pi}{2}}^{\pi} \int_0^{2\pi} a \chi(r) u_n |_{\Omega_{\text{int}}}(\mathbf{0}) \frac{a_0^{\text{ext}}}{a_0^{\text{int}} + a_0^{\text{ext}}} \frac{\alpha}{r} \partial_r u_n r^2 \sin(\theta) d\theta d\varphi \right] (r = \eta) \\
&\quad - \int_{\Omega_{\text{int}} \setminus B_\eta} a \nabla \left( \chi(r) u_n |_{\Omega_{\text{int}}}(\mathbf{0}) \frac{a_0^{\text{ext}}}{a_0^{\text{int}} + a_0^{\text{ext}}} \frac{\alpha}{r} \right) \nabla u_n r^2 \sin(\theta) dr d\theta d\varphi, \quad (3.27)
\end{aligned}$$

with the spherical coordinates  $(r, \theta, \varphi)$  defined by  $x = r \sin(\theta) \cos(\varphi)$ ,  $y = r \sin(\theta) \sin(\varphi)$  and  $z = r \cos(\theta)$ . Inserting (3.26), and (3.27) in (3.25), we obtain  $\chi(\eta) = 1$  and  $\partial_r \chi(\eta) = 0$ , for small  $\eta$

$$\begin{aligned}
\lambda_n^1 \int_{\Omega_{\text{int}}} b(u_n)^2 &= - \lim_{\eta \rightarrow 0} \left\{ \int_{\frac{\pi}{2}}^{\pi} \int_0^{2\pi} a(\mathbf{x}) \partial_r \left( u_n |_{\Omega_{\text{int}}}(\mathbf{0}) \frac{a_0^{\text{ext}}}{a_0^{\text{int}} + a_0^{\text{ext}}} \frac{\alpha}{r} \right) u_n(\mathbf{x}) r^2 \sin(\theta) d\theta d\varphi \right. \\
&\quad \left. - \int_{\frac{\pi}{2}}^{\pi} \int_0^{2\pi} a(\mathbf{x}) u_n |_{\Omega_{\text{int}}}(\mathbf{0}) \frac{a_0^{\text{ext}}}{a_0^{\text{int}} + a_0^{\text{ext}}} \frac{\alpha}{r} \partial_r u_n(\mathbf{x}) r^2 \sin(\theta) d\theta d\varphi \right\} (r = \eta). \quad (3.28)
\end{aligned}$$

Using the first order Taylor expansions of  $u_n$ ,  $\partial_r u_n$  in  $\Omega_{\text{int}}$  and (1.6), we obtain

$$u_n(\mathbf{x}) = u_n |_{\Omega_{\text{int}}}(\mathbf{0}) + O(r), \quad \partial_r u_n(\mathbf{x}) = O(1) \quad \text{and} \quad a(\mathbf{x}) = a_0^{\text{int}} + O(r). \quad (3.29)$$

Inserting (3.29) in (3.28), we have

$$\lambda_n^1 \int_{\Omega_{\text{int}}} b(u_n)^2 = \lim_{\eta \rightarrow 0} \left\{ a_0^{\text{int}} \int_{\frac{\pi}{2}}^{\pi} \int_0^{2\pi} (u_n |_{\Omega_{\text{int}}}(\mathbf{0}))^2 \frac{a_0^{\text{ext}}}{a_0^{\text{int}} + a_0^{\text{ext}}} \alpha \sin(\theta) d\theta d\varphi + O(\eta) \right\}. \quad (3.30)$$

Taking the limit, get (3.18).  $\square$

## 4 Numerical simulations

In this section, we will present two series of numeral experiments illustrating Theorem 1.1. The numerical computations are based on the parallel version of the CESC library of CERFACS (boundary element and finite element code) and of the ARPACK library (large scale eigenvalue problems solver).

For both series of experiments,  $\lambda_n^\delta$ ,  $\lambda_n$  and  $u_n$  are evaluated with a  $P_1$ -continuous (piecewise linear continuous approximation on a tetrahedral mesh) finite element approximation. The parameter  $\delta$  takes 11 values going from 1 to  $10^{-2}$ :  $\delta = 10^{-\frac{k}{5}}$  with  $0 \leq k \leq 10$ . The coefficient  $\alpha = \frac{1}{4\pi} \int_{\Sigma} \lambda_*(\mathbf{X}') d\mathbf{X}'$  is either numerically computed by solving (3.12) or approximated by  $\alpha_{\text{app}} = \frac{2}{\pi} \sqrt{\frac{A}{\pi}}$ . The numerical computation of  $\alpha$  relies on  $P_1$ -continuous (piecewise linear continuous approximation on a triangular mesh) boundary

element approximation. The computation of  $\lambda_n^\delta$  requires a refined mesh in the neighborhood of the hole. Even if the geometry is simple, 2.5 million degrees of freedom are required for the smallest  $\delta$ . The computation of  $u_n$  and  $\lambda_n$  is achieved on a coarse mesh and is therefore less costly and easier to handle (one does not have to face some errors of the mesher: For very small mesh step our mesher was simply not able to generate a regular mesh without dividing the computational domain in many regions).

We report in Figures 5, 6, 7, 8 the results of our simulations for  $n = 1$  to 4. For  $n = 0$ , we do not show the results since  $\lambda_0^\delta = \lambda_0 = \lambda_0^1 = 0$ .

During all the computations, we have tried to diminish as much as possible our numerical errors. For the smallest values of  $\delta$  the encountered linear systems become rather large (millions of unknowns) and the errors committed by the eigenvalue solver ARPACK cannot be completely neglected.

#### 4.1 First experiment

Let  $\ell_x^{\text{int}} = .6$ ,  $\ell_x^{\text{ext}} = 1.$ ,  $\ell_y = .8$ ,  $\ell_z = .3$ . The two cavities, see Figure 3, are defined by

$$\Omega_{\text{int}} = ] -\frac{\ell_x^{\text{int}}}{2}, \frac{\ell_x^{\text{int}}}{2}[\times] -\frac{\ell_y}{3}, \frac{2\ell_y}{3}[\times] -\ell_z, 0[ \text{ and } \Omega_{\text{ext}} = ] -\frac{\ell_x^{\text{ext}}}{4}, \frac{3\ell_x^{\text{ext}}}{4}[\times] -\frac{\ell_y}{3}, \frac{2\ell_y}{3}[\times] 0, \ell_z[.$$

The shape of the hole  $\Sigma$ , see Figure 4 is a polygon with vertexes  $A = (0,0)$ ,  $B = (.1,0)$ ,  $C = (.1, -.08)$ ,  $D = (-.08, -.08)$ ,  $E = (-.08, .1)$  and  $F = (0, .1)$  with

$$\alpha = 0.0578... \quad \text{and} \quad \alpha_{\text{app}} = 0.0538... \quad (4.1)$$

The coefficient functions  $a$  and  $b$  are constant and equal to 1.

#### 4.2 Second experiment

Let  $\ell_x = 1.$ ,  $\ell_y = .8$ ,  $\ell_z = .3$ . The interior cavity is

$$\Omega_{\text{int}} = ] -\frac{\ell_x}{2}, \frac{\ell_x}{2}[\times] -\frac{\ell_y}{3}, \frac{2\ell_y}{3}[\times] -\ell_z, 0[.$$

The exterior cavity is a pyramid with basis the polygon linking  $(.2, -.4, 0)$ ,  $(-.7, -.4, 0)$ ,  $(-.5, .2, 0)$  and  $(.2, .2)$  and with upper vertex  $(.1, .1, .7)$ . The shape of the hole is a centered circle of radius .1. It corresponds to

$$\alpha = \alpha_{\text{app}} = 0.0637... \quad (4.2)$$

The functions  $a$  and  $b$  are piecewise constant and given by

$$a|_{\Omega_{\text{int}}} = 2, \quad a|_{\Omega_{\text{ext}}} = 1, \quad b|_{\Omega_{\text{int}}} = 1, \quad b|_{\Omega_{\text{ext}}} = 2. \quad (4.3)$$



Figure 3: The geometry for both experiments

Figure 4: The shape of the holes  $\Sigma$  for both experiments

## 5 Conclusion

In this article we derived the first order asymptotic expansion of the eigenvalues and eigenvectors of a three dimensional elliptic operator equipped with Neumann boundary condition. This expansion allows to compute with a small computation cost a numerical approximation of these eigenvalues.

The reader can also remark that this work can easily be adapted to deal with a multiperforated straight structure if one can manage the boundary homogenisation, see [6, 17, 19]. It would be of interest to see the impact of varying coefficients on the blockage coefficient  $C$  which measures the permeability of the wall and is related to the effective size of the

$n$	1 <sup>st</sup> experiment		2 <sup>nd</sup> experiment	
	$\lambda_n$	$\lambda_n^1$	$\lambda_n$	$\lambda_n^1$
0	0	0	0	0
1	0	2.02	0	2.17
2	9.87	0.757	9.24	0.426
3	15.42	0	16.7	0.251
4	15.42	1.01	19.7	1.11

Table 1: The values of  $\lambda_n$  and  $\lambda_n^1$  (see (1.12))




Figure 5: Error  $\lambda_n^\delta - (\lambda_n + \delta\lambda_1^n)$  in log-log scale for the first experiment




Figure 6: Error  $\lambda_n^\delta - (\lambda_n + \delta\lambda_1^n)$  in log-log scale for the first experiment with  $\alpha$  replaced by  $\alpha_{app}$

hole  $s = \alpha\delta$  of Remark 3.3 and to the area  $A$  of the cell containing one hole by

$$C = \frac{A}{2\pi s}.$$

## 6 Acknowledgments.

This work was supported by the French National Research Agency under grant no. ANR-08-SYSC-001.

## References

- [1] C. AMROUCHE, V. GIRAULT, J. GIROIRE, *Dirichlet and Neumann exterior problems for the  $n$ -dimensional Laplace operator: an approach in weighted Sobolev spaces*, J. Math. Pures Appl. (9) 76 (1997), no. 1, 55–81.
- [2] A. BENDALI, A. HUARD, A. TIZAOU, S. TORDEUX AND J. P. VILA, *Asymptotic Expansions of the Eigenvalues of a 2-D boundary-value problem relative to two cavities linked by a hole of small size*, C. R. Acad. Sci. Paris, Mathématiques, 347 (2009), pp. 1147-1152.
- [3] A. BENDALI, A. TIZAOU, S. TORDEUX AND J. P. VILA, *Matching of Asymptotic Expansions for an eigenvalue problem with two cavities linked by a narrow hole*, SAM Research Report, ETH Zurich, (2009), no. 17, 74 pages.




Figure 7: Error  $\lambda_n^\delta - (\lambda_n + \delta\lambda_n^1)$  in log-log scale for the second experiment

- [4] V. BONNAILLIE-NOËL AND M. DAUGE, *Asymptotics for the low-lying eigenstates of the Schrödinger operator with magnetic field near corners*, Ann. Henri Poincaré, 7 (2006), pp. 899–931.
- [5] R. M. BROWN, P. D. HISLOP AND A. MARTINEZ, *Lower bounds on the interaction between cavities connected by a thin tube*, Duke Math. J., 73 (1994), pp. 163–176.
- [6] G. A. CHECHKIN, R. R. GADYL'SHIN, , *On boundary-value problems for the Laplacian in bounded and in unbounded domains with perforated boundaries*, Journal of Differential Equations, 216, (2005), no. 2, pp. 502–522.
- [7] M. COSTABEL, M. DAUGE, S. NICAISE, *Corner Singularities and Analytic Regularity for Linear Elliptic Systems*, in preparation.
- [8] M. DAUGE, I. DJURDJEVIC, E. FAOU, AND A. RÖSSLE, *Eigenmode asymptotics in thin elastic plates*, J. Math. Pures Appl. (9), 78 (1999), pp. 925–964.
- [9] R. DAUTRAY, J-L. LIONS, *Mathematical analysis and numerical methods for science and technology. Vol. 4. Integral equations and numerical methods*. Springer-Verlag, Berlin, 1990. x+465 pp.
- [10] R. R. GADYL'SHIN, *Surface potentials and the method of matching asymptotic expansions in the Helmholtz resonator problem*, (Russian) Algebra i Analiz 4 (1992), no. 2, 88–115, translation in St. Petersburg Math. J. 4 (1993), no. 2, 273–296.
- [11] A. M. IL'IN, *Matching of asymptotic expansions of solutions of boundary value problems*, vol. 102 of Translations of Mathematical Monographs, American Mathematical Society, Providence, RI, 1992, Translated from the Russian by V. Minachin.
- [12] V. A. KONDRATEV. *Boundary-value problems for elliptic equations in domains with conical or angular points*, Trans. Moscow Math. Soc. 16 (1967) pp. 227–313.
- [13] J. RAUCH AND M. TAYLOR, *Electrostatic screening*, J. Mathematical Phys., 16 (1975), pp. 284–288.
- [14] P.-A. RAVIART AND J.-M. THOMAS, *Introduction à l'analyse numérique des équations aux dérivées partielles*, Collection Mathématiques Appliquées pour la Maîtrise. [Collection of Applied Mathematics for the Master's Degree], Masson, Paris, (1983).
- [15] J. W. RAYLEIGH, B. STRUTT, *The Theory of Sound*, 2d ed. Dover Publications, New York, N. Y., 1945. Vol. I, xlii+480 pp.; vol. II, xii+504 pp.
- [16] J. W. RAYLEIGH, *The Theory of the Helmholtz Resonator*, Proceedings of the Royal Society of London. Series A, Containing Papers of a Mathematical and Physical Character, Vol. 92, No. 638 (Feb. 1, 1916), pp. 265–275
- [17] J. SANCHEZ-HUBERT AND E. SÁNCHEZ-PALENCIA, *Acoustic fluid flow through holes and permeability of perforated walls*, J. Math. Anal. Appl. 87 (1982), pp. 427–453.






Figure 8: The eigenvalue  $\lambda_n^\delta$ , its limit  $\lambda_n$  and its approximation  $\lambda_n + \delta\lambda_n^1$  with respect to  $\delta$

- [18] A. TAFLOV, K. UMASHANKER, B. BECKER, F. HARFOUSH, AND K. S. YEE, *Detailed ftd analysis of electromagnetic fields penetrating narrow slots and lapped joints in thin conducting screens*, IEEE Trans Antenna and Propagation, 36 (1988), pp. 247-257.
- [19] E. O. TUCK, *Matching problems involving flow through small holes*, in Advances in applied mechanics, Vol. 15, Academic Press, New York, (1975), pp. 89-158.
- [20] M. VAN DYKE, *Perturbation methods in fluid mechanics*, The Parabolic Press, Stanford, Calif., (1975).